

Propagation of partial randomness

Kojiro Higuchi

Department of Mathematics and Informatics
Faculty of Science, Chiba University
1-33 Yayoi-cho, Inage, Chiba, JAPAN
khiguchi@g.math.s.chiba-u.ac.jp

W. M. Phillip Hudelson
One Oxford Center, Suite 2600
301 Grant Street
Pittsburgh, PA 15219, USA
phil.hudelson@gmail.com

Stephen G. Simpson
Department of Mathematics
Pennsylvania State University
University Park, PA 16802, USA
<http://www.personal.psu.edu/t20>
t20@psu.edu

Keita Yokoyama
School of Information Science
Japan Advanced Institute of Science and Technology
1-1 Asahidai, Nomi, Ishikawa, 923-1292, JAPAN
y-keita@jaist.ac.jp

First draft: April 21, 2011

This draft: May 21, 2014

Abstract

Let f be a computable function from finite sequences of 0's and 1's to real numbers. We prove that strong f -randomness implies strong f -randomness relative to a PA-degree. We also prove: if X is strongly f -random and Turing reducible to Y where Y is Martin-Löf random relative to Z , then X is strongly f -random relative to Z . In addition, we prove analogous propagation results for other notions of partial randomness, including non-K-triviality and autocomplexity. We prove that f -randomness relative to a PA-degree implies strong f -randomness, hence f -randomness does not imply f -randomness relative to a PA-degree.

Keywords: partial randomness, effective Hausdorff dimension, Martin-Löf randomness, Kolmogorov complexity, models of arithmetic.

2010 Mathematics Subject Classification: Primary 03D32, Secondary 03D28, 68Q30, 03H15, 03C62, 03F30.

A version of this paper has been published in *Annals of Pure and Applied Logic*, 165(2):742–758, 2014.

Contents

Abstract	1
1 Introduction	2
2 f-randomness and strong f-randomness	4
3 g-randomness implies strong f-randomness	7
4 Propagation of strong f-randomness	8
5 Propagation of non-K-triviality	11
6 Propagation of diagonal nonrecursiveness	12
7 Propagation of autocomplexity	15
8 Vehement f-randomness	18
9 Propagation of vehement f-randomness	21
10 Other characterizations of strong f-randomness	22
11 Non-propagation of f-randomness	24
References	24

1 Introduction

We begin by recalling two known results concerning Martin-Löf randomness relative to a Turing oracle. Let \mathbb{N} denote the set of positive integers. Let $\{0, 1\}^{\mathbb{N}}$ denote the Cantor space, i.e., the set of infinite sequences of 0's and 1's.

Theorem 1.1. Let $X \in \{0, 1\}^{\mathbb{N}}$ be Martin-Löf random. Suppose X is Turing reducible to Y where Y is Martin-Löf random relative to Z . Then X is Martin-Löf random relative to Z .

Theorem 1.2. Let Q be a nonempty Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$. If $X \in \{0, 1\}^{\mathbb{N}}$ is Martin-Löf random, then X is Martin-Löf random relative to some $Z \in Q$.

Recall from [18] that a PA-degree is defined to be the Turing degree of a complete consistent theory extending first-order Peano arithmetic. It is well known that, via Gödel numbering, the set of complete consistent theories extending first-order Peano arithmetic may be viewed as a Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$. Moreover [18, 29, 32, 33], this particular Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$ is *universal* in the following sense: a Turing oracle Z is of PA-degree if and only if every nonempty Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$ contains an element which is Turing reducible to Z . Consequently, Theorem 1.2 may be restated as follows:

Theorem 1.3. If $X \in \{0, 1\}^{\mathbb{N}}$ is Martin-Löf random, then X is Martin-Löf random relative to some PA-degree.

Theorem 1.1, which we call the *XYZ Theorem*, is due to Miller/Yu [22, Theorem 4.3]. Theorems 1.2 and 1.3 are due independently to several groups of researchers: Downey et al [10, Proposition 7.4], Reimann/Slaman [27, Theorem 4.5] (also cited in [10]), and Simpson/Yokoyama [37, Lemma 3.3].

Theorems 1.2 and 1.3 have been very useful in the study of randomness. Reimann/Slaman [27] used Theorem 1.2 to prove that any noncomputable $X \in \{0, 1\}^{\mathbb{N}}$ is nonatomically random with respect to some probability measure on $\{0, 1\}^{\mathbb{N}}$. Simpson/Yokoyama [37] used a generalization of Theorem 1.2 to study the reverse mathematics of Loeb measure. Recently Brattka/Miller/Nies [5] used Theorem 1.2 to prove that $x \in [0, 1]$ is random if and only if every computable continuous function of bounded variation is differentiable at x .

The theme of Theorems 1.1–1.3 is what might be called “propagation of Martin-Löf randomness.” Namely, all of these theorems assert that if X is Martin-Löf random then X is Martin-Löf random relative to certain Turing oracles.

The purpose of this paper is to present some new results which are generalizations of Theorems 1.1–1.3. The theme of our new results might be called “propagation of partial randomness.” Here “partial randomness” refers to certain properties which are in the same vein as Martin-Löf randomness. Recent studies of partial randomness include [6, 16, 19, 21, 25, 28, 39]. Our main new results involve a specific notion of partial randomness known as *strong f -randomness* where f is an arbitrary computable function from finite sequences of 0’s and 1’s to real numbers. Along the way we present some old and new characterizations of strong f -randomness. We also consider other notions of partial randomness including *complexity* [4, 15, 19], *autocomplexity* [19], and *non- K -triviality* [11, 23].

The plan of this paper is as follows. In §2 we define *f -randomness* and *strong f -randomness* and characterize these notions in terms of Kolmogorov complexity. In §3 we prove that $(f + 2 \log_2 f)$ -randomness implies strong f -randomness. Note that §2 and §3 and §8 are largely expository. In §4 we present our main results concerning propagation of strong f -randomness. Namely, we prove appropriate generalizations of Theorems 1.1–1.3 with Martin-Löf randomness replaced by strong f -randomness. In §5 and §6 we prove analogous results concerning propagation of non- K -triviality and propagation of diagonal nonrecursiveness, respectively. In §7 we prove analogous results concerning propagation of autocomplexity, and we characterize autocomplexity in terms of f -randomness and strong f -randomness. In §8 we define *vehement f -randomness* and prove that it is equivalent to strong f -randomness, provided f is *convex*. In §9 we prove a version of Theorem 1.2 with Martin-Löf randomness replaced by vehement f -randomness. In §10 we present two new characterizations of strong f -randomness. In §11 we show that our results concerning propagation of strong f -randomness fail for f -randomness.

2 f -randomness and strong f -randomness

Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be an arbitrary computable function from finite sequences of 0's and 1's to the extended¹ real numbers. In this section we define what it means for $X \in \{0, 1\}^{\mathbb{N}}$ to be f -random, strongly f -random, f -complex, and strongly f -complex.

Recall that according to Schnorr's Theorem (see [11, Theorem 6.2.3] or [23, Theorem 3.2.9] or [34, Theorem 10.7]), X is Martin-Löf random if and only if for all n the prefix-free Kolmogorov complexity of the first n bits of X is at least n modulo an additive constant. In this section we prove generalizations of Schnorr's Theorem, replacing Martin-Löf randomness by f -randomness and strong f -randomness. Our proofs are modeled on one of the standard proofs [34, Theorem 10.7] of Schnorr's Theorem.

This section is mostly expository. For the history of the concepts and results in this section, see Calude/Staiger/Terwijn [6] and Tadaki [39].

Definition 2.1. For $X \in \{0, 1\}^{\mathbb{N}}$ and $n \in \mathbb{N}$ we write $X \upharpoonright n = X \upharpoonright \{1, \dots, n\}$ = the first n bits of X . Given $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ we define X to be f -complex or strongly f -complex if

$$\exists c \forall n (\text{KP}(X \upharpoonright n) \geq f(X \upharpoonright n) - c)$$

or

$$\exists c \forall n (\text{KA}(X \upharpoonright n) \geq f(X \upharpoonright n) - c)$$

respectively. Here KP and KA denote prefix-free complexity (see [11, §3.5] or [23, §2.2] or [34, §10]) and a priori complexity (see [11, §3.16] or [40]) respectively.

Definition 2.2. Given $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$, the f -weight of $\sigma \in \{0, 1\}^*$ is defined as $\text{wt}_f(\sigma) = 2^{-f(\sigma)}$. The direct f -weight of $A \subseteq \{0, 1\}^*$ is defined as $\text{dwt}_f(A) = \sum_{\sigma \in A} \text{wt}_f(\sigma)$. A set $P \subseteq \{0, 1\}^*$ is said to be prefix-free if no element of P is a proper initial segment of an element of P . The prefix-free f -weight of A is defined as

$$\text{pwt}_f(A) = \sup\{\text{dwt}_f(P) \mid P \subseteq A \text{ is prefix-free}\}.$$

Definition 2.3. For $\sigma \in \{0, 1\}^*$ we write $[[\sigma]] = \{X \in \{0, 1\}^{\mathbb{N}} \mid \sigma \subset X\}$. For $A \subseteq \{0, 1\}^*$ we write $[[A]] = \bigcup_{\sigma \in A} [[\sigma]]$ and

$$\widehat{A} = \{\sigma \in A \mid \nexists \rho (\rho \subset \sigma \text{ and } \rho \in A)\} = \text{the set of minimal elements of } A.$$

Note that \widehat{A} is prefix-free and $[[\widehat{A}]] = [[A]]$.

We write r.e. as an abbreviation for *recursively enumerable*. A sequence of sets $A_i \subseteq \{0, 1\}^*$, $i \in \mathbb{N}$ is said to be uniformly r.e. if $\{(\sigma, i) \mid \sigma \in A_i\}$ is r.e.

¹We define $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ to be computable if $f/(|f| + 1) : \{0, 1\}^* \rightarrow [-1, 1]$ is computable.

Definition 2.4. Assume that $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ is computable. We define $X \in \{0, 1\}^{\mathbb{N}}$ to be f -random or strongly f -random if $X \notin \bigcap_i \llbracket A_i \rrbracket$ whenever A_i is uniformly r.e. with $\text{dwt}_f(A_i) \leq 2^{-i}$ or $\text{pwt}_f(A_i) \leq 2^{-i}$ respectively.

Remark 2.5. Since $\text{pwt}_f(A) \leq \text{dwt}_f(A)$ for all A , it is clear that strong f -randomness implies f -randomness. Similarly, since $\exists c \forall \tau (\text{KA}(\tau) \leq \text{KP}(\tau) + c)$, it is clear that strong f -complexity implies f -complexity. Note also that wt_f is a *premeasure* in the sense of [26, Definition 1].

The next theorem is a straightforward generalization of Tadaki [39, Theorem 3.1].

Theorem 2.6. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. Then f -randomness is equivalent to f -complexity.

Proof. Suppose X is f -random. Let $S_i = \{\tau \mid \text{KP}(\tau) < f(\tau) - i\}$. Clearly S_i is uniformly r.e., and by Kraft's Inequality [34, Theorem 10.3] we have

$$\text{dwt}_f(S_i) = \sum_{\tau \in S_i} 2^{-f(\tau)} \leq \sum_{\tau \in S_i} 2^{-\text{KP}(\tau) - i} = 2^{-i} \sum_{\tau \in S_i} 2^{-\text{KP}(\tau)} < 2^{-i}$$

so S_i is a test for f -randomness. Since X is f -random it follows that $X \notin \bigcap_i \llbracket S_i \rrbracket$, i.e., $\exists i \forall n (\text{KP}(X \upharpoonright n) \geq f(X \upharpoonright n) - i)$, i.e., X is f -complex.

Now suppose X is not f -random, say $X \in \bigcap_i \llbracket A_i \rrbracket$ where A_i is uniformly r.e. and $\text{dwt}_f(A_i) \leq 2^{-i}$. Then

$$\sum_i \sum_{\tau \in A_{2i}} 2^{-f(\tau) + i} = \sum_i 2^i \text{dwt}_f(A_{2i}) \leq \sum_i 2^i 2^{-2i} = \sum_i 2^{-i} = 1$$

so by the Kraft/Chaitin Lemma (see [34, Corollary 10.6]) we have

$$\exists c \forall i \forall \tau (\tau \in A_{2i} \Rightarrow \text{KP}(\tau) \leq f(\tau) - i + c).$$

Since $X \in \bigcap_i \llbracket A_{2i} \rrbracket$ it follows that $\exists c \forall i \exists n (\text{KP}(X \upharpoonright n) \leq f(X \upharpoonright n) - i + c)$. In other words, X is not f -complex. This completes the proof. \square

Corollary 2.7. The sets $S_i = \{\tau \mid \text{KP}(\tau) < f(\tau) - i\}$ form a universal test for f -randomness.

Proof. Paraphrasing Theorem 2.6 we see that X is f -random if and only if $X \notin \bigcap_i \llbracket S_i \rrbracket$. It remains to prove that $\text{dwt}_f(S_i) \leq 2^{-i}$, but we have already seen this as part of the proof of Theorem 2.6. \square

The next theorem is a straightforward generalization of Calude/Staiger/Terwijn [6, Corollary 4.10].

Theorem 2.8. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. Then strong f -randomness is equivalent to strong f -complexity.

Proof. Recall that $\text{KA}(\tau) = -\log_2 m(\tau)$ where $m : \{0, 1\}^* \rightarrow [0, 1]$ is a universal left-r.e. semimeasure. See for instance [11, §3.16] or [40].

Suppose X is strongly f -random. Let $S_i = \{\tau \mid \text{KA}(\tau) < f(\tau) - i\}$. Clearly S_i is uniformly r.e. We claim that $\text{pwt}_f(S_i) \leq 2^{-i}$. To see this, let $P \subseteq S_i$ be prefix-free. Then

$$\text{dwt}_f(P) = \sum_{\tau \in P} 2^{-f(\tau)} \leq \sum_{\tau \in P} 2^{-i - \text{KA}(\tau)} = 2^{-i} \sum_{\tau \in P} m(\tau) \leq 2^{-i}$$

since m is a semimeasure. This proves our claim. Thus S_i is a test for strong f -randomness. Since X is strongly f -random, we have $X \notin \bigcap_i \llbracket S_i \rrbracket$, i.e., $\exists i \forall n (\text{KA}(X \upharpoonright n) \geq f(X \upharpoonright n) - i)$, i.e., X is strongly f -complex.

Now suppose X is not strongly f -random, say $X \in \bigcap_i \llbracket A_i \rrbracket$ where A_i is uniformly r.e. and $\text{pwt}_f(A_i) \leq 2^{-i}$. For each i let m_i be the uniformly left-r.e. semimeasure given by $m_i(\sigma) = \text{pwt}_f(\{\tau \in A_i \mid \tau \supseteq \sigma\})$. Note that $m_i(\tau) \geq \text{wt}_f(\tau)$ whenever $\tau \in A_i$. For each i we have $m_i(\langle \rangle) = \text{pwt}_f(A_i) \leq 2^{-i}$, hence $2^i m_{2i}(\langle \rangle) \leq 2^i 2^{-2i} = 2^{-i}$, so consider the left-r.e. semimeasure $\overline{m}(\sigma) = \sum_i 2^i m_{2i}(\sigma)$. Since m is a universal left-r.e. semimeasure, let c be such that $\overline{m}(\sigma) \leq 2^c m(\sigma)$ for all σ . Then for all $\tau \in A_{2i}$ we have $2^{i-f(\tau)} = 2^i \text{wt}_f(\tau) \leq 2^i m_{2i}(\tau) \leq \overline{m}(\tau) \leq 2^c m(\tau) = 2^{c-\text{KA}(\tau)}$, hence $\text{KA}(\tau) \leq f(\tau) - i + c$. Since $X \in \bigcap_i \llbracket A_{2i} \rrbracket$ it follows that $\forall i \exists n (\text{KA}(X \upharpoonright n) \leq f(X \upharpoonright n) - i + c)$. In other words, X is not strongly f -complex. This completes the proof. \square

Corollary 2.9. The sets $S_i = \{\tau \mid \text{KA}(\tau) < f(\tau) - i\}$ form a universal test for strong f -randomness.

Proof. Paraphrasing Theorem 2.8 we see that X is strongly f -random if and only if $X \notin \bigcap_i \llbracket S_i \rrbracket$. It remains to prove that $\text{pwt}_f(S_i) \leq 2^{-i}$, but we have already seen this as part of the proof of Theorem 2.8. \square

Remark 2.10. As a special case, consider the functions $f_s : \{0, 1\}^* \rightarrow [0, \infty)$ given by $f_s(\sigma) = s|\sigma|$ where s is rational and $0 < s \leq 1$. Here we are writing $|\sigma|$ = the length of σ . Define $X \in \{0, 1\}^{\mathbb{N}}$ to be s -random if it is f_s -random, and strongly s -random if it is strongly f_s -random. Note that Martin-Löf randomness is equivalent to 1-randomness and to strong 1-randomness. The effective Hausdorff dimension of X is

$$\text{effdim}(X) = \sup\{s \mid X \text{ is } s\text{-random}\} = \sup\{s \mid X \text{ is strongly } s\text{-random}\}$$

and this notion has been studied in [21, 25, 39] and many other publications.

Remark 2.11. Given a computable function $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$, it is easy to see that $\{X \mid X \text{ is } f\text{-random}\}$ and $\{X \mid X \text{ is strongly } f\text{-random}\}$ are Σ_2^0 subsets of $\{0, 1\}^{\mathbb{N}}$. Conversely, given a Σ_2^0 set $S \subseteq \{0, 1\}^{\mathbb{N}}$, we can easily construct a computable function $f : \{0, 1\}^* \rightarrow \mathbb{N}$ such that

$$S = \{X \mid X \text{ is } f\text{-random}\} = \{X \mid X \text{ is strongly } f\text{-random}\}.$$

Namely, if $S = \bigcup_i \{\text{paths through } T_i\}$ where $T_i \subseteq \{0, 1\}^*$, $i \in \mathbb{N}$ is a computable sequence of computable trees, let

$$f(\tau) = \begin{cases} 1 & \text{if } h(\tau \upharpoonright (|\tau| - 1)) = h(\tau), \\ 2|\tau| & \text{otherwise,} \end{cases}$$

where $h(\tau) =$ the least i such that $i = |\tau|$ or $\tau \in T_i$. We mention these examples in order to suggest how our concepts of f -randomness and strong f -randomness may apply to a wide variety of situations. See also Theorem 7.3 below.

3 g -randomness implies strong f -randomness

Suppose we have two computable functions $f, g : \{0, 1\}^* \rightarrow [-\infty, \infty]$. Clearly g -randomness implies f -randomness provided $\forall \sigma (f(\sigma) \leq g(\sigma))$. We now prove that g -randomness implies strong f -randomness provided g grows significantly faster than f . Our result here is a slight refinement of known results due to Calude/Staiger/Terwijn [6] and Reimann/Stephan [28]. See also Uspensky/Shen [40, §4.2].

Definition 3.1. The increasing set of $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ is

$$I(f) = \{\sigma \mid (\forall \rho \subset \sigma) (f(\rho) < f(\sigma))\}.$$

Lemma 3.2. Given a computable function $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$, we can effectively find a computable function $\bar{f} : \{0, 1\}^* \rightarrow \mathbb{N}$ such that for all σ ,

$$f_0(\sigma) < \bar{f}(\sigma) < f_0(\sigma) + 2 \quad (1)$$

where $f_0(\sigma) = \min(\max(f(\sigma), 0), 2|\sigma|)$. It then follows that f -randomness is equivalent to \bar{f} -randomness, and strong f -randomness is equivalent to strong \bar{f} -randomness.

Proof. Given $\sigma \in \{0, 1\}^*$ we can effectively approximate $f_0(\sigma)$ to find $\bar{f}(\sigma) \in \mathbb{N}$ such that (1) holds. In this way we obtain a computable function $\bar{f} : \{0, 1\}^* \rightarrow \mathbb{N}$. Using the fact that $\exists c \forall \sigma (0 < \text{KP}(\sigma) < 2|\sigma| + c$ and $0 < \text{KA}(\sigma) < 2|\sigma| + c$), we can easily see that (strong) f -complexity is equivalent to (strong) \bar{f} -complexity. The desired conclusions then follow in view of Theorems 2.6 and 2.8. \square

Lemma 3.3. Let $f : \{0, 1\}^* \rightarrow \mathbb{N}$ be computable. Given an r.e. set $A \subseteq \{0, 1\}^*$ we can effectively find an r.e. set $\bar{A} \subseteq I(f)$ such that $\llbracket A \rrbracket \subseteq \llbracket \bar{A} \rrbracket$ and $\text{dwt}_f(\bar{A}) \leq \text{dwt}_f(A)$ and $\text{pwt}_f(\bar{A}) \leq \text{pwt}_f(A)$.

Proof. Let $\bar{A} = \{\bar{\sigma} \mid \sigma \in A\}$ where $\bar{\sigma} = \min\{\rho \subseteq \sigma \mid f(\rho) \geq f(\sigma)\}$. It is straightforward to verify that this \bar{A} has the desired properties. \square

Remark 3.4. Because of Lemmas 3.2 and 3.3, we are often safe in assuming that $f : \{0, 1\}^* \rightarrow \mathbb{N}$ and that $A \subseteq I(f)$.

Theorem 3.5. Let $f, g : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable with g of the form $g(\sigma) = f(\sigma) + h(f(\sigma))$ where h is nondecreasing and $\sum_{n=1}^{\infty} 2^{-h(n)} < \infty$. If X is g -random, then X is strongly f -random.

Proof. Because h is nondecreasing, we may safely apply Lemma 3.2 to assume that $f : \{0, 1\}^* \rightarrow \mathbb{N}$. Fix c such that $\sum_n 2^{-h(n)} \leq 2^c < \infty$. Suppose X is not strongly f -random, say $X \in \bigcap_i \llbracket A_i \rrbracket$ where A_i is uniformly r.e. and $\text{pwt}_f(A_i) \leq 2^{-i}$. By Lemma 3.3 we may safely assume that $A_i \subseteq I(f)$ for all i . Let $P_{in} = \{\sigma \in A_i \mid f(\sigma) = n\}$. Clearly $A_i = \bigcup_n P_{in}$ and P_{in} is prefix-free. Thus $\text{dwt}_f(P_{in}) \leq \text{pwt}_f(A_i)$ and

$$\begin{aligned} \text{dwt}_g(A_i) &= \sum_{\sigma \in A_i} 2^{-g(\sigma)} \\ &= \sum_{\sigma \in A_i} 2^{-h(f(\sigma))} 2^{-f(\sigma)} \\ &= \sum_n \sum_{\sigma \in P_{in}} 2^{-h(n)} 2^{-f(\sigma)} \\ &= \sum_n 2^{-h(n)} \sum_{\sigma \in P_{in}} 2^{-f(\sigma)} \\ &= \sum_n 2^{-h(n)} \text{dwt}_f(P_{in}) \\ &\leq 2^c \text{pwt}_f(A_i) \\ &\leq 2^{c-i}. \end{aligned}$$

Since $X \in \bigcap_i \llbracket A_i \rrbracket$ it follows that X is not g -random, Q.E.D. \square

Theorem 3.6. Let $f : \{0, 1\}^* \rightarrow (0, \infty]$ be computable. Suppose X is $(f + (1 + \epsilon) \log_2 f)$ -random for some $\epsilon > 0$. Then X is strongly f -random.

Proof. We may safely assume that ϵ is rational. In this case it suffices to apply Theorem 3.5 with $h(x) = (1 + \epsilon) \log_2 x$. \square

Remark 3.7. Consider the computable function $f = f_s$ where $s = 1/2$, i.e., $f(\sigma) = |\sigma|/2$ for all σ . (More generally, let f be computable and satisfy certain other conditions which we shall not specify here.) Reimann/Stephan [28] have constructed an X which is f -random but not strongly f -random. Hudelson [16] has constructed an X which is strongly f -random but such that no Y Turing reducible to X is $(f + (1 + \epsilon) \log_2 f)$ -random for any $\epsilon > 0$. We conjecture that there exists an X which is f -random but such that no Y Turing reducible to X is strongly f -random.

Remark 3.8. In Theorem 3.6 and Remark 3.7 we may replace $f + (1 + \epsilon) \log_2 f$ by $f + \log_2 f + (1 + \epsilon) \log_2 \log_2 f$, etc., as in [40, §4.2].

4 Propagation of strong f -randomness

The purpose of this section is to prove generalizations of Theorems 1.1–1.3 in which Martin-Löf randomness is replaced by strong f -randomness. These generalizations are perhaps the most important new results of this paper. Let μ be the fair-coin probability measure on $\{0, 1\}^{\mathbb{N}}$ given by $\mu(\llbracket \sigma \rrbracket) = 2^{-|\sigma|}$.

Definition 4.1. A Levin system is an indexed family of sets $V_\sigma \subseteq \{0, 1\}^{\mathbb{N}}$, $\sigma \in \{0, 1\}^*$, such that

1. V_σ is Σ_1^0 uniformly in σ ,
2. $V_\sigma \supseteq V_{\sigma \frown \langle 0 \rangle} \cup V_{\sigma \frown \langle 1 \rangle}$ for all σ ,
3. $V_{\sigma \frown \langle 0 \rangle} \cap V_{\sigma \frown \langle 1 \rangle} = \emptyset$ for all σ .

These properties easily imply

4. $V_\rho \supseteq V_\sigma$ whenever $\rho \subseteq \sigma$,
5. $V_\sigma \cap V_\tau = \emptyset$ whenever σ and τ are incompatible.

Lemma 4.2. Let V_σ be a Levin system, and let f be computable. If X is strongly f -random, then $\exists c \forall n (\mu(V_{X \upharpoonright n}) \leq 2^{c-f(X \upharpoonright n)})$.

Proof. Let $A_i = \{\sigma \mid \mu(V_\sigma) > 2^{i-f(\sigma)}\}$. Clearly A_i is uniformly r.e. We claim that $\text{pwt}_f(A_i) \leq 2^{-i}$. To see this, let $P \subseteq A_i$ be prefix-free. By part 5 of Definition 4.1 we have $1 \geq \mu(\bigcup_{\sigma \in P} V_\sigma) = \sum_{\sigma \in P} \mu(V_\sigma) \geq \sum_{\sigma \in P} 2^{i-f(\sigma)} = 2^i \text{dwt}_f(P)$, so $\text{dwt}_f(P) \leq 2^{-i}$. This proves our claim. Thus A_i is a test for strong f -randomness. Since X is strongly f -random, it follows that $X \notin \llbracket A_i \rrbracket$ for some i . In other words, $\mu(V_{X \upharpoonright n}) \leq 2^{i-f(X \upharpoonright n)}$ for all n , Q.E.D. \square

Remark 4.3. Our idea of using strong f -randomness in Lemma 4.2 was inspired by Reimann's use of strong f -randomness in [26, Theorem 14].

Lemma 4.4. Let $r_\sigma, \sigma \in \{0, 1\}^*$, be a uniformly left-r.e. system of real numbers. Given a Levin system V_σ , we can effectively find a Levin system \tilde{V}_σ such that

1. $\tilde{V}_\sigma \subseteq V_\sigma$ for all σ ,
2. $\mu(\tilde{V}_\sigma) \leq r_\sigma$ for all σ ,
3. $\tilde{V}_\sigma = V_\sigma$ whenever σ is such that $\mu(V_\rho) < r_\rho$ for all $\rho \subseteq \sigma$.

Proof. The proof is awkward but straightforward. \square

Theorem 4.5. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. Suppose X is strongly f -random and Turing reducible to Y where Y is Martin-Löf random relative to Z . Then X is strongly f -random relative to Z .

Proof. Let Φ be a partial recursive functional such that $X = \Phi^Y$. Consider the Levin system $V_\sigma = \{\bar{Y} \mid \Phi^{\bar{Y}} \supseteq \sigma\}$. By Lemma 4.2 let c be such that $\mu(V_{X \upharpoonright n}) < 2^{c-f(X \upharpoonright n)}$ for all n . Applying Lemma 4.4 with $r_\sigma = 2^{c-f(\sigma)}$ we obtain a Levin system \tilde{V}_σ such that $\mu(\tilde{V}_\sigma) \leq 2^{c-f(\sigma)}$ for all σ , and $Y \in V_{X \upharpoonright n} = \tilde{V}_{X \upharpoonright n}$ for all n . Now suppose X is not strongly f -random relative to Z , say $X \in \bigcap_i \llbracket A_i^Z \rrbracket$ where A_i^Z is uniformly Z -r.e. and $\text{pwt}_f(A_i^Z) \leq 2^{-i}$. Let $W_i^Z = \bigcup_{\sigma \in A_i^Z} \tilde{V}_\sigma$. Clearly W_i^Z is uniformly $\Sigma_1^{0,Z}$. Because $X \in \bigcap_i \llbracket A_i^Z \rrbracket$ and $Y \in \bigcap_n V_{X \upharpoonright n} = \bigcap_n \tilde{V}_{X \upharpoonright n}$, we

have $Y \in \bigcap_i W_i^Z$. Let $P_i = \widehat{A}_i^Z = \{\text{minimal elements of } A_i^Z\}$. Because \widetilde{V}_σ is a Levin system, we have $W_i^Z = \bigcup_{\sigma \in P_i} \widetilde{V}_\sigma$ and hence

$$\mu(W_i^Z) = \sum_{\sigma \in P_i} \mu(\widetilde{V}_\sigma) \leq \sum_{\sigma \in P_i} 2^{c-f(\sigma)} = 2^c \text{dwt}_f(P_i) \leq 2^c \text{pwt}_f(A_i^Z) \leq 2^{c-i}$$

since P_i is a prefix-free subset of A_i^Z . Thus Y is not Martin-Löf random relative to Z , Q.E.D. \square

Remark 4.6. In Theorem 4.5 the assumption “ Y is Martin-Löf random relative to Z ” cannot be weakened to “ Y is strongly f -random relative to Z .” For example, define $Z(n) = Y(2n)$ where Y is Martin-Löf random. Then Z is strongly $1/2$ -random (indeed Martin-Löf random) and Turing reducible to Y , and Y is strongly $1/2$ -random relative to Z , but of course Z is not strongly $1/2$ -random relative to Z .

Theorem 4.7. For each $i \in \mathbb{N}$ let $f_i : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable and let $X_i \in \{0, 1\}^{\mathbb{N}}$. Suppose $\forall i (X_i \text{ is strongly } f_i\text{-random})$. Then, we can find Z of PA-degree such that $\forall i (X_i \text{ is strongly } f_i\text{-random relative to } Z)$.

Proof. By the Kučera/Gács Theorem (see [11, Theorem 8.3.2] or [23, §3.3] or [34, Theorem 3.8]), let Y be Martin-Löf random such that $\forall i (X_i \text{ is Turing reducible to } Y)$. By Theorem 1.3 let Z be of PA-degree such that Y is Martin-Löf random relative to Z . If $\forall i (X_i \text{ is strongly } f_i\text{-random})$, it follows by Theorem 4.5 that $\forall i (X_i \text{ is strongly } f_i\text{-random relative to } Z)$. \square

Corollary 4.8. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. If X is strongly f -random, then X is strongly f -random relative to some PA-degree.

Proof. Apply Theorem 4.7 with $X_i = X$ and $f_i = f$ for all i . \square

Even the following corollary appears to be new.

Corollary 4.9. Suppose $(\forall i \in \mathbb{N}) (X_i \text{ is Martin-Löf random})$. Then, we can find Z of PA-degree such that $(\forall i \in \mathbb{N}) (X_i \text{ is Martin-Löf random relative to } Z)$.

Proof. Consider $f : \{0, 1\}^* \rightarrow [0, \infty)$ where $f(\sigma) = |\sigma|$ for all σ . By Remark 2.10 X_i is Martin-Löf random if and only if X_i is strongly f -random, and similarly X_i is Martin-Löf random relative to Z if and only if X_i is strongly f -random relative to Z . Apply Theorem 4.7 with $f_i = f$ for all i . \square

We end this section by presenting a kind of Borel/Cantelli Lemma for strong f -randomness. Let us say that X is strongly BC- f -random if $\{i \mid X \in [A_i]\}$ is finite whenever A_i is uniformly r.e. and $\sum_i \text{pwt}_f(A_i) < \infty$. This notion resembles a generalization of Tadaki’s earlier notion of Solovay D -randomness [39, Definition 3.8].

Theorem 4.10. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. Suppose X is strongly f -random and Turing reducible to Y where Y is Martin-Löf random relative to Z . Then X is strongly BC- f -random relative to Z .

Proof. Suppose X is not strongly BC- f -random relative to Z . Let A_i^Z be uniformly Z -r.e. such that $\sum_i \text{pwt}_f(A_i^Z) < \infty$ and $X \in \llbracket A_i^Z \rrbracket$ for infinitely many i . Let $V_\sigma, c, \tilde{V}_\sigma, W_i^Z, P_i$ be as in the proof of Theorem 4.5. For all i we have $\mu(W_i^Z) \leq 2^c \text{pwt}_f(A_i^Z)$, hence $\sum_i \mu(W_i^Z) \leq 2^c \sum_i \text{pwt}_f(A_i^Z) < \infty$. On the other hand, for all i such that $X \in \llbracket A_i^Z \rrbracket$ we have $Y \in W_i^Z$, so $Y \in W_i^Z$ for infinitely many i . Relativizing Solovay's Lemma [34, Lemma 3.5] to Z , we see that Y is not Martin-Löf random relative to Z , Q.E.D. \square

Theorem 4.11. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. If X is strongly f -random, then X is strongly BC- f -random relative to some PA-degree.

Proof. By the Kučera/Gács Theorem, let Y be Martin-Löf random such that X is Turing reducible to Y . By Theorem 1.3 let Z be of PA-degree such that Y is Martin-Löf random relative to Z . If X is strongly f -random, Theorem 4.10 tells us that X is strongly BC- f -random relative to Z , Q.E.D. \square

Corollary 4.12. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. Then strong f -randomness is equivalent to strong BC- f -randomness.

Proof. Trivially strong BC- f -randomness implies strong f -randomness. The converse is immediate from Theorem 4.11. \square

Remark 4.13. It is possible to give a direct proof of Corollary 4.12 resembling the standard proof of Solovay's Lemma [34, Lemma 3.5].

5 Propagation of non-K-triviality

Recall from [11, 23] that X is LR-reducible to Z , abbreviated $X \leq_{\text{LR}} Z$, if $\forall Y ((Y \text{ Martin-Löf random relative to } Z) \Rightarrow (Y \text{ Martin-Löf random relative to } X))$. The concept of LR-reducibility has been very useful [20, 34, 35] in the reverse mathematics of measure-theoretic regularity. It is also known (see [11, Chapter 11] or [23, Chapter 5]) that LR-reducibility can be used to characterize K-triviality. Namely, X is K-trivial if and only if $X \leq_{\text{LR}} 0$.

From our point of view in this paper, it seems reasonable to view non-K-triviality as a kind of partial randomness notion. Accordingly, we now present appropriate analogs of our main propagation results, Theorems 4.5 and 4.7. Our results in this section are easy consequences of previously known characterizations of K-triviality.

Theorem 5.1. Suppose X is Turing reducible to Y where Y is Martin-Löf random relative to Z . Then $X \not\leq_{\text{LR}} 0$ implies $X \not\leq_{\text{LR}} Z$.

Proof. Since $X \not\leq_{\text{LR}} 0$, it follows by [11, Chapter 11] or [23, Chapter 5] that X is not a *base for Martin-Löf randomness*. In particular, since X is Turing reducible to Y , Y is not Martin-Löf random relative to X . But then, since Y is Martin-Löf random relative to Z , we have $X \not\leq_{\text{LR}} Z$, Q.E.D. \square

Theorem 5.2. Suppose $X_i \not\leq_{\text{LR}} 0$ for all $i \in \mathbb{N}$. Then, we can find Z of PA-degree such that $X_i \not\leq_{\text{LR}} Z$ for all $i \in \mathbb{N}$.

Proof. For each i let Y_i be Martin-Löf random but not Martin-Löf random relative to X_i . By Corollary 4.9 let Z be of PA-degree such that $\forall i (Y_i \text{ is Martin-Löf random relative to } Z)$. It follows that $\forall i (X_i \not\leq_{\text{LR}} Z)$, Q.E.D. \square

Remark 5.3. In Theorems 5.1 and 5.2, the conclusion $X \not\leq_{\text{LR}} Z$ implies that $X \oplus Z \not\leq_{\text{LR}} Z$, i.e., X is not K-trivial relative to Z . On the other hand, results such as Theorems 1.3 and 4.7 and 5.2 bear an obvious resemblance to the well known *GKT Theorem* (see Gandy/Kreisel/Tait [13] or [18, Theorem 2.5] or [31, Theorem VIII.2.24]). Indeed, Theorem 5.2 is just the GKT Theorem with Turing reducibility replaced by LR-reducibility.

6 Propagation of diagonal nonrecursiveness

Let $\{n\}$ denote the partial recursive functional with index n . Let DNR be the set of functions $f : \mathbb{N} \rightarrow \mathbb{N}$ which are diagonally nonrecursive, i.e., $f(n) \neq \{n\}(n)$ for all n . Known results concerning diagonal nonrecursiveness may be found in [1, 17, 19, 32]. We also consider relative DNR-ness: $\text{DNR}^Z = \{f \in \mathbb{N}^{\mathbb{N}} \mid \forall n (f(n) \neq \{n\}^Z(n))\}$. The purpose of this section is to obtain propagation results for diagonal nonrecursiveness.

Theorem 6.1. Suppose there exists a DNR function which is Turing reducible to X . Suppose also that X is Turing reducible to Y where Y is Martin-Löf random relative to Z . Then there exists a DNR^Z function which is Turing reducible to X .

In order to prove Theorem 6.1 we need the following lemma, which is a variant of the Parametrized Recursion Theorem. In stating and proving our lemma, we shall use standard recursion-theoretic notation. In particular, for any expression E we write $E \downarrow$ to mean that E is defined, and $E \uparrow$ to mean that E is undefined. We also write $E_1 \simeq E_2$ to mean that either $(E_1 \downarrow \text{ and } E_2 \downarrow \text{ and } E_1 = E_2)$ or $(E_1 \uparrow \text{ and } E_2 \uparrow)$. Via Gödel numbering, we identify finite sequences of positive integers with positive integers. We write $f \leq_{\text{T}} X$ to mean that f is Turing reducible to X .

Lemma 6.2. Let $\Theta(n, j, \sigma, -)$ be a partial recursive functional. Then, we can find a primitive recursive function $p(n, j)$ such that

$$\{p(n, j)\}(-) \simeq \Theta(n, j, \langle p(n, i) \mid i \leq j \rangle, -)$$

for all $n, j, -$.

Proof. By the Parametrized Recursion Theorem, let q be a primitive recursive function such that $\{q(n, j, \sigma)\}(-) \simeq \Theta(n, j, \sigma \hat{\ } \langle q(n, j, \sigma) \rangle, -)$ for all $n, j, \sigma, -$.

Define p primitive recursively by letting $p(n, j) = q(n, i, \langle p(n, i) \mid i < j \rangle)$ for all n, j . We then have

$$\begin{aligned}
\{p(n, j)\}(-) &\simeq \{q(n, j, \langle p(n, i) \mid i < j \rangle)\}(-) \\
&\simeq \Theta(n, j, \langle p(n, i) \mid i < j \rangle \wedge \langle q(n, j, \langle p(n, i) \mid i < j \rangle) \rangle, -) \\
&\simeq \Theta(n, j, \langle p(n, i) \mid i < j \rangle \wedge \langle p(n, j) \rangle, -) \\
&\simeq \Theta(n, j, \langle p(n, i) \mid i \leq j \rangle, -)
\end{aligned}$$

and this proves our lemma. \square

We now prove Theorem 6.1.

Proof of Theorem 6.1. Let $f \in \mathbb{N}^{\mathbb{N}}$ be DNR and $\leq_T X$. Then $f \leq_T Y$ so let Φ be a partial recursive functional such that $f = \Phi^Y$, i.e., $f(n) = \Phi(Y, n)$ for all n . As in §4 let μ be the fair-coin probability measure on $\{0, 1\}^{\mathbb{N}}$. Define a partial recursive function $\theta(n, j, \sigma) \simeq$ some m such that $\mu(\{\bar{Y} \mid \Phi^{\bar{Y}}(\sigma(j)) \downarrow = m \text{ and } (\forall i < j) (\Phi^{\bar{Y}}(\sigma(i)) \downarrow \neq \{\sigma(i)\}(\sigma(i)) \downarrow)\}) > 2^{-n}$. Apply Lemma 6.2 to obtain a primitive recursive function $p(n, j)$ such that $\{p(n, j)\}(p(n, j)) \simeq$ some m such that $\mu(V_{n, j, m}) > 2^{-n}$ where $V_{n, j, m} = \{\bar{Y} \mid \Phi^{\bar{Y}}(p(n, j)) \downarrow = m \text{ and } (\forall i < j) (\Phi^{\bar{Y}}(p(n, i)) \downarrow \neq \{p(n, i)\}(p(n, i)) \downarrow)\}$. Thus $\{p(n, j)\}(p(n, j)) \downarrow$ implies $\mu(V_{n, j}) > 2^{-n}$ where $V_{n, j} = V_{n, j, \{p(n, j)\}(p(n, j))}$. On the other hand, $i \neq j$ implies $V_{n, i} \cap V_{n, j} = \emptyset$ so for each n there is at least one $j \leq 2^n$ such that $\{p(n, j)\}(p(n, j)) \uparrow$.

Let Ψ be a partial recursive functional defined by

$$\Psi^{\bar{Y}}(n) \simeq \langle \Phi^{\bar{Y}}(p(n, i)) \mid i \leq 2^n \rangle$$

for all n . In particular we have $g \in \mathbb{N}^{\mathbb{N}}$ defined by

$$g(n) = \Psi^Y(n) = \langle f(p(n, i)) \mid i \leq 2^n \rangle$$

for all n . Clearly $g \leq_T f \leq_T X$, so it will suffice to prove that $g(n) \neq \{n\}^Z(n)$ for all but finitely many n .

Let $U_n^Z = \{\bar{Y} \mid \Psi^{\bar{Y}}(n) \downarrow = \{n\}^Z(n) \downarrow\}$. Clearly U_n^Z is uniformly $\Sigma_1^{0, Z}$. Given a rational number r , let $U_n^Z[r]$ be U_n^Z enumerated so long as its μ -measure is $\leq r$. Thus $U_n^Z[r]$ is uniformly $\Sigma_1^{0, Z}$ and $\mu(U_n^Z[r]) \leq r$. Moreover, $U_n^Z[r] = U_n^Z$ if and only if $\mu(U_n^Z) \leq r$. Since Y is Martin-Löf random, it follows by Solovay's Lemma [34, Lemma 3.5] that $Y \notin U_n^Z[2^{-n}]$ for all but finitely many n . Therefore, it will suffice to prove $g(n) \neq \{n\}^Z(n)$ for all such n .

Supposing otherwise, we would have $\Psi^Y(n) = g(n) = \{n\}^Z(n)$, hence $Y \in U_n^Z$, hence $\mu(U_n^Z) > 2^{-n}$. Moreover, for all $\bar{Y} \in U_n^Z$ we would have $\Psi^{\bar{Y}}(n) = \{n\}^Z(n) = g(n)$, hence $\Phi^{\bar{Y}}(p(n, i)) = f(p(n, i)) \neq \{p(n, i)\}(p(n, i))$ for all $i \leq 2^n$. Let $j \leq 2^n$ be such that $(\forall i < j) (\{p(n, i)\}(p(n, i)) \downarrow)$. Then $U_n^Z \subseteq V_{n, j, f(p(n, j))}$, hence $\mu(V_{n, j, f(p(n, j))}) \geq \mu(U_n^Z) > 2^{-n}$, hence $\{p(n, j)\}(p(n, j)) \downarrow$, so by induction on j we see that $\{p(n, j)\}(p(n, j)) \downarrow$ holds for all $j \leq 2^n$. This contradiction completes the proof. \square

Theorem 6.3. Let Q be a nonempty Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$. If $(\forall i \in \mathbb{N})(\exists f \in \text{DNR})(f \leq_T X_i)$, then $(\exists Z \in Q)(\forall i \in \mathbb{N})(\exists g \in \text{DNR}^Z)(g \leq_T X_i)$.

Proof. By the Kučera/Gács Theorem (see [11, Theorem 8.3.2] or [23, §3.3] or [34, Theorem 3.8]), let Y be Martin-Löf random such that $\forall i(X_i \leq_T Y)$. By Theorem 1.2 let $Z \in Q$ be such that Y is Martin-Löf random relative to Z . If $\forall i \exists f(f \in \text{DNR} \text{ and } f \leq_T X_i)$, it follows by Theorem 6.1 that $\forall i \exists g(g \in \text{DNR}^Z \text{ and } g \leq_T X_i)$. \square

Corollary 6.4. Let Q be a nonempty Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$. If there exists a DNR function which is Turing reducible to X , then for some $Z \in Q$ there exists a DNR^Z function which is Turing reducible to X .

Proof. This is the special case of Theorem 6.3 with $X_i = X$ for all $i \in \mathbb{N}$. \square

Theorem 6.5. Suppose $(\forall i \in \mathbb{N})(\exists f \in \text{DNR})(f \leq_T X_i)$. Then, there exists Z of PA-degree such that $(\forall i \in \mathbb{N})(\exists g \in \text{DNR}^Z)(g \leq_T X_i)$.

Proof. In Theorem 6.3 let Q be the Π_1^0 set consisting of all completions of first-order Peano arithmetic. \square

Corollary 6.6. If there exists a DNR function which is Turing reducible to X , then for some Z of PA-degree there exists a DNR^Z function which is Turing reducible to X .

Proof. In Corollary 6.4 let Q be the Π_1^0 set consisting of all completions of first-order Peano arithmetic. \square

Remark 6.7. As in [32, §10] and [36, §2.2], let C be a “nice” class of recursive functions. For example, C could be the class of *all* recursive functions, or the class of primitive recursive functions, or the class of recursive functions up to level α of the transfinite Ackermann hierarchy for some limit ordinal $\alpha \leq \varepsilon_0$. A function $f : \mathbb{N} \rightarrow \mathbb{N}$ is said to be C -bounded if $(\exists F \in C) \forall n(f(n) < F(n))$. In particular, f is recursively bounded if it is C -bounded where $C =$ the class of all recursive functions. Our proofs above show that Theorems 6.1 and 6.3 and 6.5 and Corollaries 6.4 and 6.6 also hold with “DNR” replaced by “ C -bounded DNR.” It suffices to note that, in our proof of Theorem 6.1, if f is C -bounded then so is g . See also the refinements mentioned in Remarks 7.8 and 7.9 below.

We end this section by presenting an alternative proof of Corollary 6.4.

Alternative proof of Corollary 6.4. Let \mathbb{N}^* be the set of finite sequences of positive integers. For each $\sigma \in \mathbb{N}^*$ let

$$Q_\sigma = \{Z \in Q \mid (\forall n < |\sigma|)(\sigma(n) \neq \{n\}^Z(n))\}$$

where $|\sigma| =$ the length of σ . Clearly $\sigma \subseteq \tau$ implies $Q_\sigma \supseteq Q_\tau$. By the Parametrization or S-m-n Theorem, let $p(n, \sigma)$ be a primitive recursive function such that for all m , $\{p(n, \sigma)\}(p(n, \sigma)) = m$ if and only if $\{n\}^Z(n) = m$ for all $Z \in Q_\sigma$. Let $f \leq_T X$ be a DNR function. Define $g \leq_T X$ recursively by

letting $g(n) = f(p(n, \langle g(i) \mid i < n \rangle))$ for all n . We are going to show that g is DNR relative to some $Z \in Q$.

We claim that $Q_{\langle g(i) \mid i < n \rangle} \neq \emptyset$ for all n . To begin with, we have $Q_{\langle \rangle} = Q \neq \emptyset$. Assume inductively that $Q_{\langle g(i) \mid i < n \rangle} \neq \emptyset$. We shall prove that $Q_{\langle g(i) \mid i \leq n \rangle} \neq \emptyset$. There are two cases. If $\{p(n, \langle g(i) \mid i < n \rangle)\}(p(n, \langle g(i) \mid i < n \rangle)) = m$, we have $\{n\}^Z(n) = m$ for all $Z \in Q_{\langle g(i) \mid i < n \rangle}$, but $g(n) = f(p(n, \langle g(i) \mid i < n \rangle)) \neq m$ since f is DNR. Thus $Q_{\langle g(i) \mid i \leq n \rangle} = Q_{\langle g(i) \mid i < n \rangle} \neq \emptyset$. If $\{p(n, \langle g(i) \mid i < n \rangle)\}(p(n, \langle g(i) \mid i < n \rangle))$ is undefined, there exists $Z \in Q_{\langle g(i) \mid i < n \rangle}$ such that $\{n\}^Z(n) \neq g(n)$, and then Z belongs to $Q_{\langle g(i) \mid i \leq n \rangle}$. This proves our claim.

By compactness, our claim implies that $\bigcap_{n=0}^{\infty} Q_{\langle g(i) \mid i < n \rangle} \neq \emptyset$. Moreover, from the definition of $Q_{\langle g(i) \mid i < n \rangle}$ we see that g is DNR relative to any $Z \in \bigcap_{n=0}^{\infty} Q_{\langle g(i) \mid i < n \rangle}$. This completes the proof. \square

Remark 6.8. Our alternative proof of Corollary 6.4 is more constructive than the previous proof via Theorem 6.1 and the Kučera/Gács Theorem. In particular, the alternative proof can be formalized in WKL_0 (see [31]) while the previous proof cannot.

There are some issues here which are interesting from the viewpoint of reverse mathematics [31]. For example, consider the following statement.

Let Q be a nonempty Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$. If X_1 and X_2 are Martin-Löf random, there exists $Z \in Q$ such that X_1 and X_2 are Martin-Löf random relative to Z .

By Corollary 4.9 this statement is true, and from the truth of the statement it follows easily that the statement is true in all ω -models of WKL_0 . Moreover, we conjecture that the statement is provable in WKL_0 . On the other hand, by [2, Theorem 2.1] together with [38], the following special case of the Kučera/Gács Theorem is false in all ω -models of WKL_0 except those which contain $0^{(1)}$ = the Turing jump of 0.

If X_1 and X_2 are Martin-Löf random, there exists a Martin-Löf random Y such that $X_1 \leq_T Y$ and $X_2 \leq_T Y$.

7 Propagation of autocomplexity

In this section we prove propagation results for *autocomplexity* and *complexity*. We also obtain a characterization of autocomplexity in terms of f -randomness and strong f -randomness. With this characterization plus [19, Theorem 2.3], we see considerable overlap between the propagation results of this section and those of §4 and §6.

Definition 7.1.

1. Following [19] we define $X \in \{0, 1\}^{\mathbb{N}}$ to be autocomplex if there exists an unbounded function $h : \mathbb{N} \rightarrow \mathbb{N}$ such that $h \leq_T X$ and $h(n) \leq \text{KS}(X \upharpoonright n)$ for all n . Here KS denotes simple Kolmogorov complexity [40], also known as plain complexity [11, 23].

2. Following [4] and [15] and [19], we define $X \in \{0, 1\}^{\mathbb{N}}$ to be complex if there exists an unbounded computable function $h : \mathbb{N} \rightarrow \mathbb{N}$ such that $h(n) \leq \text{KS}(X \upharpoonright n)$ for all n .

Remark 7.2. By [40, §4.3.1] there exist constants c_1 and c_2 such that $\text{KS}(\sigma) \leq \text{KP}(\sigma) + c_1 \leq \text{KS}(\sigma) + 3 \log_2 |\sigma| + c_2$ and $\text{KA}(\sigma) \leq \text{KP}(\sigma) + c_1 \leq \text{KA}(\sigma) + 3 \log_2 |\sigma| + c_2$ for all $\sigma \in \{0, 1\}^*$. These inequalities imply that the distinctions among KS and KP and KA are immaterial for some purposes. In particular, we can replace KS in Definition 7.1 by KP or KA.

We begin with autocomplexity.

Theorem 7.3. The following are pairwise equivalent.

1. X is autocomplex.
2. X is f -random for some computable $f : \{0, 1\}^* \rightarrow \mathbb{N}$ such that $\{f(X \upharpoonright n) \mid n \in \mathbb{N}\}$ is unbounded.
3. X is strongly f -random for some computable $f : \{0, 1\}^* \rightarrow \mathbb{N}$ such that $\{f(X \upharpoonright n) \mid n \in \mathbb{N}\}$ is unbounded.

Proof. The equivalence $2 \Leftrightarrow 3$ is clear in view of Remark 7.2.

To prove $2 \Rightarrow 1$, suppose 2 holds via f . By Theorem 2.6 X is f -complex, so let $c \in \mathbb{N}$ be such that $\text{KP}(X \upharpoonright n) \geq f(X \upharpoonright n) - c$ for all n . Then for all n we have $\text{KP}(X \upharpoonright n) \geq h(n)$ where $h(n) = \max(1, f(X \upharpoonright n) - c)$. Clearly $h \leq_{\text{T}} X$ (in fact h is Lipschitz computable from X) and h is unbounded, so it follows by Remark 7.2 that X is autocomplex, i.e., 1 holds.

It remains to prove $1 \Rightarrow 2$. Suppose X is autocomplex. By Remark 7.2 let $h : \mathbb{N} \rightarrow \mathbb{N}$ be unbounded such that $h \leq_{\text{T}} X$ and $h(n) \leq \text{KP}(X \upharpoonright n)$ for all n . Let Φ be a partial recursive functional such that $h = \Phi^X$. Consider the primitive recursive function $f : \{0, 1\}^* \rightarrow \mathbb{N}$ defined by $f(\sigma) = \max\{p(\sigma, n) \mid n \leq |\sigma|\}$ where $p(\sigma, n) = \Phi_{|\sigma|}^{\sigma}(n)$ if $\Phi_{|\sigma|}^{\sigma}(n) \downarrow$, otherwise $p(\sigma, n) = 1$. Then for all n and all sufficiently large $m \geq n$ we have $h(n) = p(X \upharpoonright m, n) \leq f(X \upharpoonright m)$. Since $\{h(n) \mid n \in \mathbb{N}\}$ is unbounded, it follows that $\{f(X \upharpoonright m) \mid m \in \mathbb{N}\}$ is unbounded. Consider the primitive recursive function $q(\sigma) =$ the least $n \leq |\sigma|$ such that $f(\sigma) = p(\sigma, n)$. Let c be a constant such that $\text{KP}(\sigma \upharpoonright q(\sigma)) \leq \text{KP}(\sigma) + c$ for all σ . Then for all m we have $\text{KP}(X \upharpoonright m) + c \geq \text{KP}(X \upharpoonright q(X \upharpoonright m)) \geq h(X \upharpoonright q(X \upharpoonright m)) \geq p(X \upharpoonright m, q(X \upharpoonright m)) = f(X \upharpoonright m)$ so X is f -complex. It follows by Theorem 2.6 that X is f -random. This completes the proof. \square

Theorem 7.4.

1. If X is autocomplex and $\leq_{\text{T}} Y$ where Y is Martin-Löf random relative to Z , then X is autocomplex relative to Z .
2. If $(\forall i \in \mathbb{N})(X_i \text{ is autocomplex})$, there exists Z of PA-degree such that $(\forall i \in \mathbb{N})(X_i \text{ is autocomplex relative to } Z)$.

First proof. Part 1 is immediate from Theorems 4.5 and 7.3. Part 2 is immediate from Theorems 4.7 and 7.3. \square

Second proof. By Kjos-Hanssen/Merkle/Stephan [19, Theorem 2.3] we know that X is autocomplex if and only if there exists a DNR function which is Turing reducible to X . Modulo this result, parts 1 and 2 are equivalent to Theorems 6.1 and 6.5 respectively. \square

Remark 7.5. Yet another proof of Theorem 7.4 was obtained independently by Bienvenu [3] who had seen it conjectured in an earlier draft of the present paper. The earlier draft included Theorems 4.5 and 4.7, as well as Corollary 6.4 with our alternative proof, but it did not include Theorem 6.1 or 6.5 or 7.3.

We now turn to propagation results for complexity. Let us define $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ to be length-invariant if $\forall \sigma \forall \tau (|\sigma| = |\tau| \Rightarrow f(\sigma) = f(\tau))$.

Theorem 7.6. The following are pairwise equivalent.

1. X is complex.
2. X is f -random for some computable $f : \{0, 1\}^* \rightarrow \mathbb{N}$ which is unbounded and length-invariant.
3. X is strongly f -random for some computable $f : \{0, 1\}^* \rightarrow \mathbb{N}$ which is unbounded and length-invariant.

Proof. This is immediate from Theorems 2.6 and 2.8 and Remark 7.2. \square

Theorem 7.7.

1. If X is complex and $\leq_T Y$ where Y is Martin-Löf random relative to Z , then X is complex relative to Z .
2. If $(\forall i \in \mathbb{N}) (X_i \text{ is complex})$, there exists Z of PA-degree such that $(\forall i \in \mathbb{N}) (X_i \text{ is complex relative to } Z)$.

Proof. Part 1 is immediate from Theorems 4.5 and 7.6. Part 2 is immediate from Theorems 4.7 and 7.6. \square

Remark 7.8. By [19, Theorem 2.3] we know that X is complex if and only if some DNR function is truth-table reducible to X . Consequently, the Turing degrees of complex X 's are the same as the Turing degrees of recursively bounded DNR functions. And of course, the Turing degrees of autocomplex X 's are the same as the Turing degrees of DNR functions. Thus Theorems 4.5 and 4.7 may be viewed as far-reaching refinements not only of Theorems 7.4 and 7.7 but also of Theorems 6.1–6.5 and Remark 6.7.

Remark 7.9. By [1, Theorem 1.8] there exists an autocomplex X such that no complex Y is Turing reducible to X . Within the class of complex X 's, much more refined results of the same kind have been obtained by Hudelson [16] generalizing the main result of Miller [21, Theorem 4.1]. See also Remarks 3.7 and 3.8 above, as well as [36, §§2.1–2.3, Figure 1].

8 Vehement f -randomness

In this section we define vehement f -randomness and discuss its relationship with strong f -randomness. The notion of vehement f -randomness was originally introduced by Kjos-Hanssen (unpublished, but see [26]). We prove that, under a convexity hypothesis on f , vehement f -randomness is equivalent to strong f -randomness. Our result is a generalization of known results due to Reimann [26, Corollary 21] and Miller [21, Lemma 3.3].

Definition 8.1. Given $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$, the vehement f -weight of $A \subseteq \{0, 1\}^*$ is defined as $\text{vwt}_f(A) = \inf\{\text{dwt}_f(S) \mid \llbracket A \rrbracket \subseteq \llbracket S \rrbracket\}$.

Remark 8.2. Note that $\llbracket A \rrbracket \subseteq \llbracket B \rrbracket$ implies $\text{vwt}_f(A) \leq \text{vwt}_f(B)$. In particular, $\text{vwt}_f(A)$ depends only on $\llbracket A \rrbracket$.

Lemma 8.3. For all A we have $\text{vwt}_f(A) \leq \text{dwt}_f(\widehat{A}) \leq \text{pwt}_f(A)$.

Proof. The first inequality holds because $\llbracket A \rrbracket \subseteq \llbracket \widehat{A} \rrbracket$. The second inequality holds because \widehat{A} is a prefix-free subset of A . \square

Definition 8.4. Fix $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$. A good cover of A is a set B such that $\llbracket A \rrbracket \subseteq \llbracket B \rrbracket$ and $\text{pwt}_f(B) \leq \text{vwt}_f(A)$. It follows by Remark 8.2 and Lemma 8.3 that $\text{vwt}_f(A) = \text{vwt}_f(B) = \text{dwt}_f(\widehat{B}) = \text{pwt}_f(B)$.

Lemma 8.5. Suppose B is a good cover of A . Given $F \subseteq \widehat{B}$ let us write $A_F = \{\sigma \in A \mid \llbracket F \rrbracket \supseteq \llbracket \sigma \rrbracket\}$ and $B_F = \{\tau \in B \mid \llbracket F \rrbracket \supseteq \llbracket \tau \rrbracket\}$. Then B_F is a good cover of A_F .

Proof. Clearly $\widehat{B}_F = F$, hence $\llbracket A_F \rrbracket \subseteq \llbracket F \rrbracket = \llbracket \widehat{B}_F \rrbracket = \llbracket B_F \rrbracket$. In order to show that B_F is a good cover of A_F , it remains to show that $\text{dwt}_f(P) \leq \text{dwt}_f(S)$ whenever $P \subseteq B_F$ is prefix-free and $\llbracket A_F \rrbracket \subseteq \llbracket S \rrbracket$. Letting $G = \widehat{B} \setminus F$ we see that $P \cap G = \emptyset$ and $P \cup G$ is a prefix-free subset of B and $\llbracket A \rrbracket = \llbracket A_F \rrbracket \cup \llbracket A_G \rrbracket \subseteq \llbracket S \rrbracket \cup \llbracket G \rrbracket = \llbracket S \cup G \rrbracket$. Thus $\text{dwt}_f(P) + \text{dwt}_f(G) = \text{dwt}_f(P \cup G) \leq \text{pwt}_f(B) \leq \text{vwt}_f(A) \leq \text{dwt}_f(S \cup G) \leq \text{dwt}_f(S) + \text{dwt}_f(G)$, hence $\text{dwt}_f(P) \leq \text{dwt}_f(S)$, Q.E.D. \square

Remark 8.6. Let B be a good cover of A , and suppose τ is such that $\llbracket B \rrbracket \not\supseteq \llbracket \tau \rrbracket$. Then obviously no initial segment of τ belongs to B . In other words, $\tau \in \widehat{B \cup \{\tau\}}$. Letting $F = \widehat{B \cup \{\tau\}} \setminus \{\tau\}$ and applying Lemma 8.5, we see that $\widehat{B}_F = F$ and B_F is a good cover of A_F .

Definition 8.7. We define $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ to be convex if $\text{wt}_f(\sigma) \leq \text{wt}_f(\sigma \frown \langle 0 \rangle) + \text{wt}_f(\sigma \frown \langle 1 \rangle)$ for all $\sigma \in \{0, 1\}^*$. Equivalently, $\text{wt}_f(\sigma) \leq \text{dwt}_f(S)$ for all $\sigma \in \{0, 1\}^*$ and all $S \subseteq \{0, 1\}^*$ such that $\llbracket S \rrbracket = \llbracket \sigma \rrbracket$.

Lemma 8.8. Assume that f is convex. Suppose B is a good cover of A but not of $A' = A \cup \{\sigma\}$. Choose $\tau \subseteq \sigma$ so as to minimize $\text{dwt}_f(\widehat{B \cup \{\tau\}})$. Then $B' = B \cup \{\tau\}$ is a good cover of A' .

Proof. Obviously $\llbracket B' \rrbracket \supseteq \llbracket A' \rrbracket$ so it remains to prove that $\text{dwt}_f(P') \leq \text{dwt}_f(S')$ whenever $P' \subseteq B'$ is prefix-free and $\llbracket A' \rrbracket \subseteq \llbracket S' \rrbracket$.

Since $\llbracket \sigma \rrbracket \subseteq \llbracket A' \rrbracket \subseteq \llbracket S' \rrbracket$, let $\tau^* \subseteq \sigma$ be as short as possible such that $\llbracket \tau^* \rrbracket \subseteq \llbracket S' \rrbracket$. Obviously S' contains no proper initial segment of τ^* . Hence $\llbracket \tau^* \rrbracket = \llbracket S^* \rrbracket$ for some $S^* \subseteq S'$. It follows by Definition 8.7 that $\text{wt}_f(\tau^*) \leq \text{dwt}_f(S^*)$. Therefore, replacing S' by $(S' \setminus S^*) \cup \{\tau^*\}$, we may safely assume that $\tau^* \in S'$.

Since $\llbracket \sigma \rrbracket \not\subseteq \llbracket B \rrbracket$ and $\tau \subseteq \sigma$ and $\tau^* \subseteq \sigma$, we obviously have $\llbracket \tau \rrbracket \not\subseteq \llbracket B \rrbracket$ and $\llbracket \tau^* \rrbracket \not\subseteq \llbracket B \rrbracket$. Applying Remark 8.6 to τ and to τ^* , we obtain sets $F = \widehat{B \cup \{\tau\}} \setminus \{\tau\}$ and $F^* = \widehat{B \cup \{\tau^*\}} \setminus \{\tau^*\}$. In particular, since $\llbracket A \rrbracket \subseteq \llbracket S' \rrbracket$ we have $\llbracket A_{F^*} \rrbracket = \llbracket A \rrbracket \setminus \llbracket \tau^* \rrbracket \subseteq \llbracket S' \rrbracket \setminus \llbracket \tau^* \rrbracket \subseteq \llbracket S' \setminus \{\tau^*\} \rrbracket$. Moreover, by our choice of τ we have $\text{dwt}_f(B \cup \{\tau\}) \leq \text{dwt}_f(B \cup \{\tau^*\})$.

We are now ready to complete the proof of Lemma 8.8. If $\tau \notin P'$ we have $P' = P \subseteq B$, hence $\text{dwt}_f(P) \leq \text{pwt}_f(B) \leq \text{vwt}_f(A) \leq \text{dwt}_f(S')$ and we are done. Suppose now that $\tau \in P'$. Then $P' = P \cup \{\tau\}$ where $P \subseteq B_F$. Thus we have

$$\begin{aligned}
\text{dwt}_f(P') &= \text{dwt}_f(P) + \text{wt}_f(\tau) \\
&\leq \text{pwt}_f(B_F) + \text{wt}_f(\tau) \\
&= \text{dwt}_f(F) + \text{wt}_f(\tau) \\
&= \text{dwt}_f(\widehat{B \cup \{\tau\}}) \\
&\leq \text{dwt}_f(\widehat{B \cup \{\tau^*\}}) \\
&= \text{dwt}_f(F^*) + \text{wt}_f(\tau^*) \\
&= \text{vwt}_f(A_{F^*}) + \text{wt}_f(\tau^*) \\
&\leq \text{dwt}_f(S' \setminus \{\tau^*\}) + \text{wt}_f(\tau^*) \\
&= \text{dwt}_f(S')
\end{aligned}$$

and again we are done. \square

Definition 8.9. Given $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ define

$$L_f : \{(P_1, P_2) \mid P_1, P_2 \text{ are finite and prefix-free}\} \rightarrow \{0, 1\}$$

by

$$L_f(P_1, P_2) = \begin{cases} 1 & \text{if } \text{dwt}_f(P_1) < \text{dwt}_f(P_2), \\ 0 & \text{otherwise.} \end{cases}$$

We say that f is strongly computable if both f and L_f are computable. This is often the case, e.g., if f is computable and integer-valued as in Lemma 3.2. Note also that Lemma 3.3 depends only on strong computability.

Lemma 8.10. Let f be strongly computable and convex. If A is r.e., we can effectively find an r.e. set B such that B is a good cover of A .

Proof. For $n = 0, 1, 2, \dots$ let A_n consist of the first n elements in some fixed recursive enumeration of A . Assume inductively that we have found a finite set B_n which is a good cover of A_n . Let $A_{n+1} = A_n \cup \{\sigma_n\}$. If $\llbracket \sigma_n \rrbracket \subseteq \llbracket B_n \rrbracket$ let $B_{n+1} = B_n$. Otherwise, use strong computability to effectively choose $\tau_n \subseteq \sigma_n$ which minimizes $\text{dwt}_f(\widehat{B_n \cup \{\tau_n\}})$. Lemma 8.8 then implies that $B_{n+1} = B_n \cup \{\tau_n\}$ is a good cover of A_{n+1} . Finally let $B = \bigcup_{n=1}^{\infty} B_n$. Clearly B is r.e. and $\llbracket A \rrbracket \subseteq \llbracket B \rrbracket$, so it remains to prove that $\text{dwt}_f(P) \leq \text{vwt}_f(A)$ for all prefix-free sets $P \subseteq B$. But clearly $\text{dwt}_f(P) = \sup\{\text{dwt}_f(P_0) \mid P_0 \text{ is a finite subset of } P\}$, so it suffices to consider finite prefix-free sets. If $P \subseteq B$ is finite and prefix-free, let n be such that $P \subseteq B_n$. Then $\text{dwt}_f(P) \leq \text{pwt}_f(B_n) \leq \text{vwt}_f(A_n) \leq \text{vwt}_f(A)$, Q.E.D. \square

Lemma 8.11. Let f be strongly computable and convex. If A is r.e., we can effectively find an r.e. set B such that $\llbracket A \rrbracket \subseteq \llbracket B \rrbracket$ and $\text{pwt}_f(B) \leq \text{vwt}_f(A)$.

Proof. This is a restatement of Lemma 8.10. \square

Definition 8.12. Assume that $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ is computable. We define $X \in \{0, 1\}^*$ to be vehemently f -random if $X \notin \bigcap_i \llbracket A_i \rrbracket$ whenever A_i is uniformly r.e. such that $\text{vwt}_f(A_i) \leq 2^{-i}$.

Theorem 8.13. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be strongly computable and convex. Then vehement f -randomness is equivalent to strong f -randomness.

Proof. Suppose X is not strongly f -random, say $X \in \bigcap_i \llbracket A_i \rrbracket$ where A_i is uniformly r.e. and $\text{pwt}_f(A_i) \leq 2^{-i}$. By Lemma 8.3 we have $\text{vwt}_f(A_i) \leq \text{pwt}_f(A_i) \leq 2^{-i}$ so X is not vehemently f -random.

Now suppose X is not vehemently f -random, say $X \in \bigcap_i \llbracket A_i \rrbracket$ where A_i is uniformly r.e. and $\text{vwt}_f(A_i) \leq 2^{-i}$. By Lemma 8.11 we can find uniformly r.e. B_i such that $\llbracket A_i \rrbracket \subseteq \llbracket B_i \rrbracket$ and $\text{pwt}_f(B_i) \leq \text{vwt}_f(A_i) \leq 2^{-i}$. Clearly $X \in \bigcap_i \llbracket B_i \rrbracket$, so X is not strongly f -random. \square

We now sketch how to replace “strongly computable” by “computable.”

Lemma 8.14. Let f be computable and convex. Given $\epsilon > 0$ we can effectively find an \bar{f} which is strongly computable and convex and such that $f(\sigma) < \bar{f}(\sigma) < f(\sigma) + \epsilon$ for all σ .

Proof. Let \mathbb{Q} be the set of rational numbers. By a straightforward but awkward construction, we can find $\bar{f} : \{0, 1\}^* \rightarrow \mathbb{Q}$ which is computable and convex and such that $f(\sigma) < \bar{f}(\sigma) < f(\sigma) + \epsilon$ for all σ . From the \mathbb{Q} -valuedness of \bar{f} it follows easily that \bar{f} is strongly computable. \square

Lemma 8.15. Let f be computable and convex. Given $\delta > 0$ and an r.e. set A , we can effectively find an r.e. set B such that $\llbracket A \rrbracket \subseteq \llbracket B \rrbracket$ and $\text{pwt}_f(B) \leq (1 + \delta) \cdot \text{vwt}_f(A)$.

Proof. Let \bar{f} be as in Lemma 8.14 with $\epsilon = \log_2(1 + \delta)$. If A is r.e., apply Lemma 8.11 to find an r.e. set B such that $\llbracket A \rrbracket \subseteq \llbracket B \rrbracket$ and $\text{pwt}_{\bar{f}}(B) \leq \text{vwt}_{\bar{f}}(A)$. It is then easy to check that $\text{pwt}_f(B) \leq (1 + \delta) \cdot \text{vwt}_f(A)$. \square

Theorem 8.16. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable and convex. Then vehement f -randomness is equivalent to strong f -randomness.

Proof. Proceed as in the proof of Theorem 8.13 but instead of Lemma 8.11 use Lemma 8.15 with $\delta = 1$. \square

9 Propagation of vehement f -randomness

In this section we present an alternative proof of one of our main results concerning propagation of strong f -randomness, Corollary 4.8. Our alternative proof proceeds via vehement f -randomness and depends heavily on Remark 8.2. Our alternative proof has the advantage of being a direct generalization of one of the known proofs (see [10, Proposition 7.4]) of the corresponding result for Martin-Löf randomness, Theorem 1.2.

Theorem 9.1. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable and convex. Let Q be a nonempty Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$. If X is strongly f -random, then X is strongly f -random relative to some $Z \in Q$.

Proof. Relativizing Corollary 2.9 let S_i^Z be a universal test for strong f -randomness relative to Z . In other words, S_i^Z is uniformly r.e. relative to Z and $\text{pwt}_f(S_i^Z) \leq 2^{-i}$ and $\forall X \forall Z (X \notin \bigcap_i \llbracket S_i^Z \rrbracket \Leftrightarrow X \text{ is strongly } f\text{-random relative to } Z)$. By Lemma 8.3 we have $\text{vwt}_f(S_i^Z) \leq \text{pwt}_f(S_i^Z) \leq 2^{-i}$ so by Theorem 8.16 S_i^Z is also a universal test for vehement f -randomness relative to Z . Thus, letting $U_i^Z = \llbracket S_i^Z \rrbracket$, we have

$$\forall X \forall Z (X \notin \bigcap_i U_i^Z \Leftrightarrow X \text{ is vehemently } f\text{-random relative to } Z)$$

and U_i^Z is uniformly Σ_1^0 relative to Z .

Let $\tilde{U}_i = \bigcap_{Z \in Q} U_i^Z$. Since Q is Π_1^0 , it follows by compactness that \tilde{U}_i is uniformly Σ_1^0 . Therefore, let \tilde{S}_i be uniformly r.e. such that $\llbracket \tilde{S}_i \rrbracket = \tilde{U}_i$. For any $Z \in Q$ we have $\tilde{U}_i \subseteq U_i^Z$, i.e., $\llbracket \tilde{S}_i \rrbracket \subseteq \llbracket S_i^Z \rrbracket$, so $\text{vwt}_f(\tilde{S}_i) \leq \text{vwt}_f(S_i^Z) \leq 2^{-i}$ by Remark 8.2. Thus \tilde{S}_i is a test for vehement f -randomness. In particular we have $\forall X (X \text{ vehemently } f\text{-random} \Rightarrow X \notin \bigcap_i \tilde{U}_i)$.

Suppose now that X is strongly f -random. By Theorem 8.16 X is vehemently f -random, so

$$X \notin \bigcap_i \tilde{U}_i = \bigcap_i \bigcap_{Z \in Q} U_i^Z = \bigcap_{Z \in Q} \bigcap_i U_i^Z.$$

Let $Z \in Q$ be such that $X \notin \bigcap_i U_i^Z$. Then X is vehemently f -random relative to Z , so by Theorem 8.16 X is strongly f -random relative to Z , Q.E.D. \square

Theorem 9.2. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable and convex. Let Q be a nonempty Π_1^0 subset of $\{0, 1\}^{\mathbb{N}}$. If X is vehemently f -random, then X is vehemently f -random relative to some $Z \in Q$.

Proof. This is immediate from Theorems 8.16 and 9.1. \square

10 Other characterizations of strong f -randomness

In this section we present two new characterizations of strong f -randomness. One of our new characterizations is in terms of f -randomness relative to a PA-degree. The other is in terms of what we call *provable noncomplexity*.

Theorem 10.1. Let $f : \{0, 1\}^* \rightarrow [-\infty, \infty]$ be computable. The following are pairwise equivalent.

1. X is strongly f -random.
2. X is strongly f -random relative to some PA-degree.
3. X is f -random relative to some PA-degree.

Proof. The implication $1 \Rightarrow 2$ follows from Theorem 4.7. The implication $2 \Rightarrow 3$ is trivial. It remains to prove $3 \Rightarrow 1$. Assume that 1 fails, i.e., X is not strongly f -random. Let A_i , $i \in \mathbb{N}$ be uniformly r.e. such that $\text{pwt}_f(A_i) \leq 2^{-i}$ and $X \in \bigcap_i \llbracket A_i \rrbracket$. For each i let \widehat{A}_i be the set of minimal elements of A_i . Let Q be the set of sequences Z_i , $i \in \mathbb{N}$ such that $Z_i \subseteq \{0, 1\}^*$ and $\text{dwt}_f(Z_i) \leq 2^{-i}$ and $\forall \sigma (\sigma \in A_i \Rightarrow \exists \rho (\rho \subseteq \sigma \text{ and } \rho \in Z_i))$. The sequence \widehat{A}_i , $i \in \mathbb{N}$ belongs to Q , so Q is nonempty. Moreover, Q may be viewed as a Π_1^0 set in the Cantor space. Therefore, given Z of PA-degree, we can find a sequence B_i , $i \in \mathbb{N}$ which is Turing reducible to Z and belongs to Q . From the definition of Q it follows that $\text{dwt}_f(B_i) \leq 2^{-i}$ and $X \in \bigcap_i \llbracket B_i \rrbracket$. Thus X is not f -random relative to Z . This holds for all PA-degrees, so 3 fails, Q.E.D. \square

For our second characterization, let PA denote first-order Peano arithmetic. Within PA we define prefix-free complexity $\text{KP} : \{0, 1\}^* \rightarrow \mathbb{N}$ and a priori complexity $\text{KA} : \{0, 1\}^* \rightarrow (0, \infty)$ as usual. Also within PA we define $\text{KP}^{(j)} =$ prefix-free complexity relative to $0^{(j)}$, and $\text{KA}^{(j)} =$ a priori complexity relative to $0^{(j)}$, where $0^{(j)} =$ the j th Turing jump of 0. Let $f : \{0, 1\}^* \rightarrow \mathbb{N}$ and $X \in \{0, 1\}^{\mathbb{N}}$ be arbitrary.

Definition 10.2. Let K stand for KP or KA, and let Z be a Turing oracle. We define X to be K^Z - f -complex if $\exists c \forall n (K^Z(X \upharpoonright n) > f(X \upharpoonright n) - c)$.

Definition 10.3. Let M be a nonstandard model of PA. We define X to be M - f -complex if $\forall r \exists c \forall n (\text{KP}_r(X \upharpoonright n) > f(X \upharpoonright n) - c)$. Here r ranges over M -finite functions r with prefix-free domain, and $\text{KP}_r(\tau) = \min\{|\sigma| \mid r(\sigma) = \tau\}$.

Definition 10.4. Let K stand for KP or KA or $\text{KP}^{(j)}$ or $\text{KA}^{(j)}$. Let T be a consistent theory extending PA. We define X to be provably K - f -noncomplex in T if $\forall c \exists n (T \vdash (\exists m < n) (K(X \upharpoonright m) < f(X \upharpoonright m) - c))$.

Theorem 10.5. Let K stand for KP or KA. Let T be a recursively axiomatizable, consistent theory extending PA. Let $f : \{0, 1\}^* \rightarrow \mathbb{N}$ and $X \in \{0, 1\}^{\mathbb{N}}$ be arbitrary. For each $j \in \mathbb{N}$ the following are pairwise equivalent.

1. X is not K^Z - f -complex for any Z of PA-degree.

2. X is not M - f -complex for any nonstandard $M \models T$.
3. X is provably K - f -noncomplex in some recursively axiomatizable, consistent theory extending T .
4. X is provably $K^{(j)}$ - f -noncomplex in some recursively axiomatizable, consistent theory extending T .

Proof. This is a special case of [41, Theorems 4.1 and 4.4]. □

Theorem 10.6. Let $f : \{0, 1\}^* \rightarrow \mathbb{N}$ be computable. Let T be a recursively axiomatizable, consistent theory extending PA. For all $X \in \{0, 1\}^{\mathbb{N}}$ the following are pairwise equivalent.

1. X is strongly f -random.
2. X is M - f -complex for some nonstandard $M \models T$.
3. X is not provably KP- f -noncomplex in any recursively axiomatizable, consistent theory extending T .

Proof. By Theorems 2.8 and 10.1 X is strongly- f -random if and only if X is KA^Z - f -complex for all Z of PA-degree. The equivalences $1 \Leftrightarrow 2$ and $1 \Leftrightarrow 3$ then follow by Theorem 10.5. □

Define K^Z -length-complexity to mean K^Z - f -complexity where $f(\sigma) = \text{length of } \sigma$, and similarly for M -length-complexity and provable KP-length-noncomplexity.

Theorem 10.7. Let T be a recursively axiomatizable, consistent theory extending PA. For all $X \in \{0, 1\}^{\mathbb{N}}$ the following are pairwise equivalent.

1. X is Martin-Löf random.
2. X is M -length-complex for some nonstandard $M \models T$.
3. X is not provably KP-length-noncomplex in any recursively axiomatizable, consistent theory extending T .

Proof. This is the special case of Theorem 10.6 with $f(\sigma) = |\sigma|$ for all σ . □

Remark 10.8. By Theorem 10.5 our notion of provable length-noncomplexity is stable under relativation to a strong oracle. Thus, letting X be Martin-Löf random and Turing reducible to $0^{(1)}$, we see that X is not $\text{KP}^{(1)}$ -length-complex but not provably $\text{KP}^{(1)}$ -noncomplex in any recursively axiomatizable, consistent extension T of PA. It follows that for any such T there exist $\tau \in \{0, 1\}^*$ and $n \in \mathbb{N}$ such that $\text{KP}^{(1)}(\tau) < n$ but $T \not\vdash \text{KP}^{(1)}(\tau) < n$. Comparing this to the celebrated Chaitin Incompleteness Theorem [7, 24], we now have a somewhat different example of a statement which is true but not provable in T .

11 Non-propagation of f -randomness

In this section we show that Theorems 4.5 and 4.7 and 10.6 fail if strong f -randomness is replaced by f -randomness.

Theorem 11.1. For many f 's, e.g., $f(\sigma) = |\sigma|/2$, we can find an X which is f -random but not f -random relative to any PA-degree.

Proof. By Reimann/Stephan [28] let X be f -random but not strongly f -random. By Theorem 10.1 X is not f -random relative to any PA-degree. \square

Corollary 11.2. For many f 's, e.g., $f(\sigma) = |\sigma|/2$, we can find an X which is f -random but provably KP- f -noncomplex in some recursively axiomatizable, consistent extension of PA. Indeed, X is provably KP- f -noncomplex in some recursively axiomatizable, consistent extension of any recursively axiomatizable, consistent extension of PA.

Proof. By Theorem 11.1 let X be f -random but not f -random relative to any PA-degree. The desired conclusion follows by Theorem 10.5. \square

References

- [1] Klaus Ambos-Spies, Bjørn Kjos-Hanssen, Steffen Lempp, and Theodore A. Slaman. Comparing DNR and WWKL. *Journal of Symbolic Logic*, 69:1089–1104, 2004. 12, 17
- [2] George Barmpalias, Andrew E. M. Lewis, and Keng Meng Ng. The importance of Π_1^0 classes in effective randomness. *Journal of Symbolic Logic*, 75(1):387–400, 2010. 15
- [3] Laurent Bienvenu. Personal communication. June 14, 2012. 17
- [4] Stephen Binns. Π_1^0 classes with complex elements. *Journal of Symbolic Logic*, 73:1341–1353, 2008. 3, 16
- [5] Vasco Brattka, Joseph S. Miller, and André Nies. Randomness and differentiability. *Transactions of the American Mathematical Society*, 2014 (estimated). Preprint, <http://arxiv.org/abs/1104.4465v4>, 39 pages; to appear in TAMS pending acceptance of revised version. 3
- [6] Cristian Calude, Ludwig Staiger, and Sebastiaan A. Terwijn. On partial randomness. *Annals of Pure and Applied Logic*, 138:20–30, 2006. 3, 4, 5, 7
- [7] Gregory J. Chaitin. Information-theoretic limitations of formal systems. *Journal of the ACM*, 21(3):403–424, 1974. 23

- [8] Z. Chatzidakis, P. Koepke, and W. Pohlers, editors. *Logic Colloquium '02: Proceedings of the Annual European Summer Meeting of the Association for Symbolic Logic and the Colloquium Logicum, held in Münster, Germany, August 3–11, 2002*. Number 27 in Lecture Notes in Logic. Association for Symbolic Logic, 2006. VIII + 359 pages. [27](#)
- [9] J. C. E. Dekker, editor. *Recursive Function Theory*. Proceedings of Symposia in Pure Mathematics. American Mathematical Society, 1962. VII + 247 pages. [26](#)
- [10] Rod Downey, Denis R. Hirschfeldt, Joseph S. Miller, and André Nies. Relativizing Chaitin’s halting probability. *Journal of Mathematical Logic*, 5:167–192, 2005. [3](#), [21](#)
- [11] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic Randomness and Complexity*. Theory and Applications of Computability. Springer, 2010. XXVIII + 855 pages. [3](#), [4](#), [6](#), [10](#), [11](#), [14](#), [15](#)
- [12] J.-E. Fenstad, I. T. Frolov, and R. Hilpinen, editors. *Logic, Methodology and Philosophy of Science VIII*. Number 126 in Studies in Logic and the Foundations of Mathematics. North-Holland, 1989. XVII + 702 pages. [25](#)
- [13] Robin O. Gandy, Georg Kreisel, and William W. Tait. Set existence. *Bulletin de l’Académie Polonaise des Sciences, Série des Sciences Mathématiques, Astronomiques et Physiques*, 8:577–582, 1960. [12](#)
- [14] S. S. Goncharov, R. Downey, and H. Ono, editors. *Mathematical Logic in Asia: Proceedings of the 9th Asian Logic Conference, Novosibirsk*. World Scientific Publishing Company, Ltd., 2006. VIII + 328 pages. [26](#)
- [15] Kojiro Higuchi and Takayuki Kihara. Effective strong nullness and effectively closed sets. In S. B. Cooper, A. Dawar, and B. Löwe, editors, *CiE 2012*, number 7318 in Lecture Notes in Computer Science, pages 303–312. Springer-Verlag, 2012. [3](#), [16](#)
- [16] W. M. Phillip Hudelson. Mass problems and initial segment complexity. *Journal of Symbolic Logic*, 79(1):20–44, 2014. [3](#), [8](#), [17](#)
- [17] Carl G. Jockusch, Jr. Degrees of functions with no fixed points. In [\[12\]](#), pages 191–201, 1989. [12](#)
- [18] Carl G. Jockusch, Jr. and Robert I. Soare. Π_1^0 classes and degrees of theories. *Transactions of the American Mathematical Society*, 173:35–56, 1972. [2](#), [12](#)
- [19] Bjørn Kjos-Hanssen, Wolfgang Merkle, and Frank Stephan. Kolmogorov complexity and the recursion theorem. *Transactions of the American Mathematical Society*, 363:5465–5480, 2011. [3](#), [12](#), [15](#), [16](#), [17](#)

- [20] Bjørn Kjos-Hanssen, Joseph S. Miller, and David Reed Solomon. Lowness notions, measure and domination. *Proceedings of the London Mathematical Society*, 85(3):869–888, 2012. [11](#)
- [21] Joseph S. Miller. Extracting information is hard: a Turing degree of non-integral effective Hausdorff dimension. *Advances in Mathematics*, 226:373–384, 2011. [3](#), [6](#), [17](#), [18](#)
- [22] Joseph S. Miller and Liang Yu. On initial segment complexity and degrees of randomness. *Transactions of the American Mathematical Society*, 360:3193–3210, 2008. [3](#)
- [23] André Nies. *Computability and Randomness*. Oxford University Press, 2009. XV + 433 pages. [3](#), [4](#), [10](#), [11](#), [14](#), [15](#)
- [24] Panu Raatikainen. On interpreting Chaitin’s Incompleteness Theorem. *Journal of Philosophical Logic*, 27(6):569–586, 1998. [23](#)
- [25] Jan Reimann. *Computability and Fractal Dimension*. PhD thesis, University of Heidelberg, 2004. XI + 130 pages. [3](#), [6](#)
- [26] Jan Reimann. Effectively closed sets of measures and randomness. *Annals of Pure and Applied Logic*, 156:170–182, 2008. [5](#), [9](#), [18](#)
- [27] Jan Reimann and Theodore A. Slaman. Measures and their random reals. *Transactions of the American Mathematical Society*, 2014 (estimated). Preprint, <http://arxiv.org/abs/0802.2705v2>, 17 pages. [3](#)
- [28] Jan Reimann and Frank Stephan. On hierarchies of randomness tests. In [\[14\]](#), pages 215–232, 2006. [3](#), [7](#), [8](#), [24](#)
- [29] Dana S. Scott. Algebras of sets binumerable in complete extensions of arithmetic. In [\[9\]](#), pages 117–121, 1962. [2](#)
- [30] S. G. Simpson, editor. *Reverse Mathematics 2001*. Number 21 in Lecture Notes in Logic. Association for Symbolic Logic, 2005. X + 401 pages. [26](#)
- [31] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Perspectives in Mathematical Logic. Springer-Verlag, 1999. XIV + 445 pages; Second Edition, Perspectives in Logic, Association for Symbolic Logic, Cambridge University Press, 2009, XVI+ 444 pages. [12](#), [15](#)
- [32] Stephen G. Simpson. Mass problems and randomness. *Bulletin of Symbolic Logic*, 11:1–27, 2005. [2](#), [12](#), [14](#)
- [33] Stephen G. Simpson. Π_1^0 sets and models of WKL_0 . In [\[30\]](#), pages 352–378, 2005. [2](#)
- [34] Stephen G. Simpson. Almost everywhere domination and superhighness. *Mathematical Logic Quarterly*, 53:462–482, 2007. [4](#), [5](#), [10](#), [11](#), [13](#), [14](#)

- [35] Stephen G. Simpson. Mass problems and measure-theoretic regularity. *Bulletin of Symbolic Logic*, 15:385–409, 2009. [11](#)
- [36] Stephen G. Simpson. Mass problems associated with effectively closed sets. *Tohoku Mathematical Journal*, 63(4):489–517, 2011. [14](#), [17](#)
- [37] Stephen G. Simpson and Keita Yokoyama. A nonstandard counterpart of WWKL. *Notre Dame Journal of Formal Logic*, 52:229–243, 2011. [3](#)
- [38] Frank Stephan. Martin-Löf random and PA-complete sets. In [\[8\]](#), pages 341–347, 2006. [15](#)
- [39] Kohtaro Tadaki. A generalization of Chaitin’s halting probability Ω and halting self-similar sets. *Hokkaido Mathematical Journal*, 31:219–253, 2002. [3](#), [4](#), [5](#), [6](#), [10](#)
- [40] V. A. Uspensky and A. Shen. Relations between varieties of Kolmogorov complexity. *Mathematical Systems Theory*, 29:271–292, 1996. [4](#), [6](#), [7](#), [8](#), [15](#), [16](#)
- [41] Keita Yokoyama. A generalization of Levin-Schnorr’s theorem. October 2013. 11 pages, <http://arxiv.org/abs/1310.3091>. [23](#)